

NDSS


National Diabetes Services Scheme

An Australian Government Initiative

NDSS Helpline 1800 637 700
ndss.com.au

Gestational Diabetes

Caring for yourself
and your baby



Find this resource at ndss.com.au

 **diabetes**
australia

The NDSS is administered by Diabetes Australia

Contents

What is gestational diabetes?	2
Who is at increased risk of gestational diabetes?	4
How is gestational diabetes diagnosed?	5
Why does gestational diabetes need to be treated?	6
Who can help you with gestational diabetes?	7
Emotional health	8
How is gestational diabetes managed?	9
Healthy eating	10
Pregnancy weight gain	21
Physical activity	22
Monitoring blood glucose levels	25
Medication	27
Hypoglycaemia	30
The birth	32
After the birth	34
Looking after your future health	37
National Diabetes Services Scheme	38

If receiving information about pregnancy or gestational diabetes raises personal concerns for you or causes you any distress, you can opt out from receiving further communications by visiting our website at ndss.com.au/gdm-update or calling the **NDSS Helpline 1800 637 700**. If you need support, ask your GP or local hospital maternity service about support services available in your area.

Disclaimer:

This information is intended as a guide only. It should not replace individual medical advice. If you have any concerns about your health, or further questions, you should contact your health professionals.

Gestational Diabetes

Caring for yourself and your baby

Having gestational diabetes is common. In Australia, one in six pregnant women will develop gestational diabetes during pregnancy.

Gestational diabetes is a form of diabetes that can develop around the 24th to 28th week of pregnancy. For most women, diabetes goes away after the baby is born.

Finding out you have gestational diabetes may come as a shock. You may be worried about how this will affect your pregnancy and your baby. While there is an increased risk of complications during pregnancy and birth, the good news is that the risks of health problems for mother and baby are reduced if gestational diabetes is carefully managed.

This booklet gives you information about gestational diabetes, looking after your health during pregnancy and where to get the support you need to manage gestational diabetes. This booklet provides general information only, it is important to discuss your individual care with your team of health professionals.

What is gestational diabetes?

Diabetes is a condition where there is too much glucose (sugar) in the bloodstream.


Glucose is an important source of energy for your body. It comes from carbohydrate foods that you eat, such as bread, pasta, rice, cereals, fruits, starchy vegetables, milk and yoghurt. Your body breaks down carbohydrates into glucose, which then enters your bloodstream.

Insulin is needed to move glucose from the bloodstream into the body's cells where it is used for energy. Insulin is a hormone made in the body by your pancreas.


During pregnancy, some of the hormones produced by the placenta (which provides nutrition for your growing baby), reduce how well your body's insulin works. This is called insulin resistance.

Insulin resistance means that the pancreas then needs to make extra insulin to keep blood glucose levels in the healthy range. If the pancreas is unable to make enough insulin, blood glucose levels rise and gestational diabetes develops.

Gestational diabetes will not lead to your baby being born with diabetes.



Glucose and insulin changes in gestational diabetes



No gestational diabetes: Insulin moves glucose from the blood into your cells.

Gestational diabetes: Insulin resistance and not enough insulin lead to less glucose entering the cells so blood glucose levels become higher.

When the baby is born, insulin requirements fall, blood glucose levels return to the normal range (for a non-pregnant woman) and diabetes usually disappears.

Who is at increased risk of gestational diabetes?

Women with an increased risk of gestational diabetes include those who:

- » have had gestational diabetes in a previous pregnancy
- » are older, especially over 40 years of age
- » are from an Aboriginal or Torres Strait Islander background
- » are from a Melanesian, Polynesian, Indian subcontinent, Middle Eastern or Asian background
- » have had elevated blood glucose levels in the past
- » have a family history of type 2 diabetes or a first-degree relative (mother or sister) who has had gestational diabetes
- » are above the healthy weight range
- » have polycystic ovary syndrome
- » have gained weight too rapidly in the first half of pregnancy
- » have had a large baby (weighing more than 4,500g) or complications in a previous pregnancy
- » are taking some types of antipsychotic or steroid medications.

Some women without known risk factors will also develop gestational diabetes.

How is gestational diabetes diagnosed?

All women should be checked for gestational diabetes during pregnancy. This usually occurs between 24 and 28 weeks of pregnancy, although some women may be advised to be checked earlier.

An oral glucose tolerance test (OGTT) is used to check how your body responds to a glucose load. After fasting (not eating) for 8–12 hours, a blood sample is taken. You then have a drink containing 75g of glucose, and blood samples are taken one and two hours later.

If your blood glucose level is above the normal range at your fasting, one or two-hour test, you have gestational diabetes.

Even if your blood glucose levels are only just above the healthy range, you still have gestational diabetes. Your health professionals will discuss your results and work with you to develop a plan for managing your gestational diabetes. Your gestational diabetes will be managed based on what best suits your needs. This will be different for each person.



Why does gestational diabetes need to be treated?

If blood glucose levels are high during pregnancy, the extra glucose passes through the placenta to the baby, who then makes extra insulin.

This can make the baby grow too big, which can cause problems during labour and increase the risk of early delivery or the need for a caesarean section.

After the birth, the baby may have a greater risk of low blood glucose levels (hypoglycaemia). This is because the baby is no longer receiving extra glucose from their mother, but they continue to make more insulin than a baby usually would, causing their blood glucose levels to drop.

Women with gestational diabetes are also at greater risk of developing high blood pressure and pre-eclampsia (high blood pressure, protein in the urine and fluid retention or swelling) during pregnancy.

Carefully managed gestational diabetes reduces the risk of these health problems for mother and baby. Your doctor and a team of diabetes health professionals can work with you to help keep your blood glucose levels within the target range to provide the best outcome for both you and your baby.

Most women with gestational diabetes will have a healthy pregnancy and a healthy baby.

Who can help you with gestational diabetes?


When you are diagnosed with gestational diabetes, there are health professionals who can help you manage this condition.

Your diabetes health care team may vary, depending on where you live or where you go for your pregnancy care and how your gestational diabetes is managed. This team may include:

- » an endocrinologist – a doctor who specialises in diabetes
- » a credentialled diabetes educator or a diabetes nurse practitioner – a specialist nurse or midwife who will help you and your family learn how to monitor and manage your blood glucose levels
- » an accredited practising dietitian – who will help you with a healthy eating plan for your pregnancy
- » your general practitioner (GP).

Your team of diabetes health professionals will work closely with your pregnancy team that may include:

- » an obstetrician – a doctor who specialises in pregnancy and birth
- » a midwife – who will provide support, care and advice during pregnancy, labour and after your baby is born
- » a physiotherapist or accredited exercise physiologist – who may help you and your partner prepare for the birth process and advise on exercise during and after pregnancy.



Emotional health

Being diagnosed with gestational diabetes may come as a shock. Your first reactions may be disbelief, sadness, anger or self-blame. At this time, it is common to feel a mixture of emotions.

There are many different risk factors for gestational diabetes, so it is important to know that getting gestational diabetes is not your fault. You may feel uncertain or concerned about how gestational diabetes will affect the health of your baby. You may also feel anxious about your blood glucose levels. Remember however, that with carefully managed gestational diabetes most women will go on to have a healthy pregnancy and healthy baby.

Some women feel overwhelmed by all of the extra appointments, the new information and the new skills needed to manage gestational diabetes. Getting the right information and support can help.

You don't have to go through this alone. Let your partner, family and friends know how you feel so they can support you. Let them know to what extent you want them to be involved in managing your gestational diabetes and invite them to share their feelings too. Your team of health professionals and the NDSS are here to support you.

If you are having problems coping with the diagnosis of gestational diabetes or feeling anxious, worried or overwhelmed, your health professionals can advise you about support services available to you locally.

In Australia, one in six women will develop gestational diabetes - you are not alone.




How is gestational diabetes managed?

You can manage gestational diabetes by:

- » following a healthy eating plan
- » doing regular physical activity
- » monitoring blood glucose levels, and
- » taking medication (if needed).

Managing gestational diabetes will help keep blood glucose levels within the target range for a healthy pregnancy. Eating well and being active will also help you to manage your pregnancy weight gain.



Healthy eating

When you have gestational diabetes, following a healthy eating plan can help to keep your blood glucose levels within the target range, provide nutrition for you and your growing baby and achieve healthy weight changes during your pregnancy.

Healthy eating for gestational diabetes includes:

- » eating 3 moderate sized meals and 2–3 snacks, spread out over the day
- » choosing the right type and amount of carbohydrate foods at each meal and snack
- » choosing foods that are low in saturated fat
- » making high-fibre food choices
- » eating a variety of foods that provide the nutrients you need during pregnancy

Nutrients required in higher amounts for pregnancy include iron (found in red meat, chicken, fish, legumes), folate (found in dark green leafy vegetables) and iodine (found in fish, bread, dairy foods).

Your dietitian can advise you on how to get the nutrition you need for you and your baby, while helping you to make healthy food choices that will help you manage your blood glucose levels.



Carbohydrates

Carbohydrate foods are broken down into glucose and used by the body for energy. They are very important for you and your baby.

Carbohydrate foods include:

- » breads and breakfast cereals
- » pasta, rice, noodles, couscous, quinoa, cracked wheat
- » starchy vegetables (such as potato, corn, sweet potato, taro and cassava)
- » fruits
- » legumes (such as baked beans, chickpeas and lentils)
- » milk and yoghurt.

To manage your blood glucose levels, you need to eat the right **amount** and **type** of carbohydrate foods.

Amount of carbohydrate

Spreading your carbohydrate foods over 3 meals and 2–3 snacks each day can help manage your blood glucose levels. Large amounts of carbohydrate foods at any one meal or snack can cause blood glucose levels to rise too high. Your dietitian can advise you on the amounts of carbohydrate that are right for you.

In some women, blood glucose levels continue to be high, even with healthy eating and regular physical activity. If this happens to you, make an appointment with your diabetes health professionals, including your dietitian, to see if any food changes may help. It is important not to cut out carbohydrates as the baby requires carbohydrate as its main energy source. Some women may need medication, usually insulin injections, to manage their blood glucose levels.



Type of carbohydrate

Carbohydrates break down into glucose at different rates. The glycemic index (GI) tells us how slowly or quickly carbohydrate foods affect blood glucose levels.

High-GI carbohydrate foods break down into glucose quickly, which means a higher and faster rise in blood glucose levels after eating.

Low-GI carbohydrate foods break down into glucose slowly. They result in a smaller and slower rise in blood glucose levels after eating.

Foods with a lower GI are a better choice when trying to manage blood glucose levels. The best carbohydrate food choices are those that are high in fibre and have a lower GI.

Lower GI carbohydrate choices include:

Breads - dense wholegrain/seeded breads, multigrain bread, raisin bread, mixed grain wraps

Cereals – traditional or steel cut rolled oats, natural muesli

Pasta – most fresh and dried pasta cooked ‘al dente’ (not overcooked)

Noodles – vermicelli, hokkien, buckwheat, udon, soba or fresh rice noodles

Rice - SunRice® Low GI rice (white/brown) or basmati rice (white/brown)


Grains – pearl barley, cracked wheat and quinoa

Dairy and alternatives – milk, soy milk and yoghurt

Vegetables and legumes – sweet potato, corn, taro, lentils, chickpeas, kidney beans

Fruit – most fresh fruit, canned fruit in natural juice

When making low GI food choices, it is still important to consider the amount of carbohydrate you eat, as recommended by your dietitian.



Acknowledgement: glycemicindex.com

Sugars and sweeteners

Sugars (including white, raw and brown sugar, honey, glucose and syrups) are also carbohydrates, but they provide no nutritional benefit and can cause your blood glucose level to rise too high.

Limit added sugars and avoid foods and drinks that are high in sugar and have little nutritional value, such as soft drinks, cordials, fruit juices, cakes, biscuits, chocolates and lollies.

There are a range of alternative sweeteners available to replace sugar. While these are not necessary, you may still choose to use them to add sweetness without adding sugar. Small amounts of alternative sweeteners can be used during pregnancy. All of the sweeteners approved for use in Australia have been tested and deemed safe by Food Standards Australia New Zealand.

Fats

Use small amounts of healthy fats, such as olive or canola oils and spreads, unsalted nuts, seeds and avocado.

Limit the amount of saturated fat you eat by choosing lean meats, skinless chicken and low-fat dairy foods. Limit butter, ghee, cream, sour cream and coconut milk/cream.

Avoid takeaways and processed foods high in saturated fats such as pastries, packaged biscuits, savoury snacks, chocolates and processed meats. If eaten in large amounts, fats can cause extra weight gain, which can make gestational diabetes harder to manage.



Protein

Protein is important for the maintenance of your body and the growth of your baby. Protein can also help you feel full for longer. Protein foods include lean meat, skinless chicken, fish, eggs, tofu and reduced-fat cheese. Milk, yoghurt and legumes (beans, lentils, chickpeas) are also important sources of protein (these foods also contain carbohydrate).

Eat plenty of vegetables

Eating plenty of vegetables is important for a healthy pregnancy. They are a good source of fibre, vitamins and minerals. Most vegetables are low in carbohydrate and will not affect your blood glucose levels (except for potato, corn, sweet potato, taro and cassava).

Eat a variety of different coloured vegetables and salads, such as tomatoes, cucumber, celery, capsicum, onions, cauliflower, zucchini, broccoli, spinach, peas, cabbage, lettuce, green beans, eggplant, carrot, leek, squash, pumpkin and Asian greens.

Talk to your dietitian about making healthy food choices for managing both your diabetes and general health during pregnancy.



Putting together a healthy meal

When putting together a healthy main meal:


- » Fill half your plate with salad (home-prepared) or vegetables (not including potato, corn, sweet potato, taro or cassava).
- » Choose a lower GI carbohydrate food such as pasta, basmati rice (white/brown), SunRice® Low GI rice (white/brown), wholegrain bread, corn, sweet potato, low-GI noodles or legumes (such as lentils, chickpeas, kidney beans).
- » Add a moderate serve of protein food, such as lean meat (beef, lamb, pork), skinless chicken, fish/seafood, tofu or eggs.

Include healthy fats and oils as part of a balanced meal, such as olive oil in cooking or as a salad dressing, avocado in a salad or nuts/seeds in a stir-fry.

Lower GI carbohydrate food such as:

- » 1 cup cooked pasta **OR**
- » 1 cup cooked low-GI rice **OR**
- » 1 cup cooked low-GI noodles **OR**
- » 1 cup sweet potato + 1 small corn cob **OR**
- » ½ cup cooked legumes + ⅔ cup cooked pasta **OR**
- » 2 slices wholegrain bread + 1 serve of fruit.

Plenty of vegetables



Moderate serve of protein food

Image acknowledgment: Diabetes SA

The sample meal ideas on the following pages provide suggested amounts of carbohydrate at each meal and snack. Your dietitian will advise you on the amount of carbohydrate to suit your individual needs.

Sample meal ideas

The following healthy meal ideas provide a starting point for planning meals and snacks to help you manage your gestational diabetes. The amounts shown here are the suggested serving sizes for one person and are a guide only.

To make sure you are eating the amounts that are right for you, ask your dietitian for advice based on your individual needs and food preferences.

As the amount of carbohydrate in food products and recipes varies, checking your blood glucose levels after meals will help you adjust food choices to suit your individual needs.

Follow pregnancy food safety guidelines as advised by your health professionals when preparing meals and snacks; see page 20.



Breakfast ideas

Choose one of the following options:

- ☐ Porridge: 1 cup cooked traditional oats served with reduced-fat milk, 1 tablespoon natural or plain Greek yoghurt and ½ cup sliced strawberries
- ☐ 1–2 slices wholegrain toast with poly or monounsaturated spread, served with 1–2 boiled eggs, plus 1 cup reduced-fat milk or soy milk (with added calcium)
- ☐ ½ cup untoasted muesli or 1 cup high fibre cereal (e.g. All-Bran Wheat Flakes® or Guardian®) with reduced-fat milk and 1 tablespoon natural or plain Greek yoghurt
- ☐ 1 cup fresh fruit salad served with ½ cup natural or plain Greek yoghurt and 30g mixed nuts/seeds
- ☐ 1–2 slices wholegrain toast lightly spread with avocado or peanut butter, plus 1 serve of whole fresh fruit
- ☐ 2–3 small idli with ¼ cup sambar, plus chopped cucumber and tomatoes
- ☐ 1 medium bowl (1½ cups) of congee (try brown/black rice or a mixture of rice, rolled oats, barley or lentils) with vegetables and egg, tofu or lean meat

Lunch ideas

Choose one of the following options:

- ☐ 1 wholegrain bread roll/sandwich/medium wrap filled with boiled egg/reduced-fat cheese/tinned salmon/tinned tuna/freshly cooked chicken with salad and avocado, plus 1 serve of whole fresh fruit
- ☐ 4 Vita-Weats™ with avocado, tomato and reduced-fat cheese, plus 100g tub fruit yoghurt and 1 serve of whole fresh fruit
- ☐ Pasta salad: 1 cup cooked pasta, tinned tuna and salad vegetables served with olive oil vinaigrette
- ☐ 2 egg and vegetable omelette served with 2 slices wholegrain toast, plus 1 serve of whole fresh fruit
- ☐ Stir-fry with a moderate serve of lean beef/chicken/fish/tofu and vegetables*, plus 1 cup cooked (small Chinese bowl) low-GI noodles
- ☐ 1 medium bowl (2 cups) homemade vegetable and lentil soup, plus 1 slice wholegrain bread
- ☐ 1 cup dhal with 1–2 small (15cm) chapati/roti (try multigrain atta or ½ chickpea flour and ½ wholemeal flour) and salad

* not including potato, sweet potato, corn, taro or cassava




Dinner ideas

Choose one of the following options:

- ☐ Serve of lean beef Bolognese sauce with 1 cup cooked pasta, served with garden salad and a drizzle of olive oil dressing
- ☐ Serve of cooked fresh fish with 1 cup cooked sweet potato, 1 small corn cob, steamed green vegetables and lemon wedges
- ☐ Stir-fry with a moderate serve of lean chicken/beef/pork/tofu and vegetables* served with 1 cup cooked (small Chinese bowl) low-GI rice (white/brown)
- ☐ Tomato-based casserole with a moderate serve of lean lamb/skinless chicken/beef and vegetables* served with 1 cup cooked pasta
- ☐ Moderate serve of pan-fried lean steak/skinless chicken/salmon with a small wholegrain roll, a garden salad and 1 small corn cob
- ☐ Moderate serve lean meat-based curry (gosht) or soya chunks (Nutrela™) curry with green vegetables served with 1 cup cooked basmati rice (white/brown) and salad
- ☐ Asian style soup (fat removed) with vegetables* and lean meat/fish/tofu/skinless chicken and 1 cup cooked low-GI noodles

* not including potato, sweet potato, corn, taro or cassava



Snack suggestions

Choose 1–2 of the following options for snacks, such as morning tea, afternoon tea and supper (before-bed snack):

- ☐ 1 serve whole fresh fruit (e.g. apple, orange, small banana or pear)
- ☐ 2–3 small fruits (e.g. mandarins, plums)
- ☐ $\frac{3}{4}$ cup canned fruit in natural juice (drained)
- ☐ 100g tub fruit yoghurt
- ☐ 1 cup reduced-fat milk or soy milk (with added calcium)
- ☐ 200g tub natural or plain Greek yoghurt topped with fresh berries or passionfruit
- ☐ One thin slice wholegrain fruit toast with poly or monounsaturated spread
- ☐ Small handful (30g) unsalted mixed nuts and seeds with 1 tablespoon sultanas or 6 dried apricot halves or 2–3 small dates
- ☐ 4 Vita-Weats™ + reduced fat cheese + vegetable sticks (carrot, cucumber, celery, capsicum)
- ☐ 2 wholegrain Ryvita® crispbread with reduced-fat cheese and tomato
- ☐ 1 x 25g packet Happy Snack Company Roasted Chickpeas™
- ☐ 1 small cob of corn or $\frac{1}{2}$ small golden sweet potato



Food safety

Pregnant women are at greater risk of food poisoning and should prepare and store food carefully. This includes preparing raw and cooked foods separately, avoiding raw or undercooked meat/chicken/seafood and following food cooking and storage instructions. Ensure all salads (including fruit salad) are home-prepared, and fruits and vegetables are washed thoroughly. Cook and reheat food until steaming hot and keep cold foods refrigerated.

Protecting yourself from exposure to high-risk foods that can cause infections and harm your developing baby is also very important. Avoid foods that may contain listeria bacteria, such as soft cheeses, unpasteurised dairy products, pre-cooked cold chicken, soft serve ice cream, deli/sandwich meats, bean sprouts, rockmelon, pre-prepared salads, tahini, hommous and pâté. Avoid raw or undercooked eggs, as these may contain salmonella.

Certain types of fish, including shark/flake, swordfish and deep-sea perch, also need to be limited during pregnancy due to their high mercury content.

Seek advice from your dietitian and/or state health department on guidelines for food safety during pregnancy.

Drinks

Drinks such as cordial, juice and soft drinks are high in kilojoules and sugar and can cause your blood glucose levels to rise too high. Choose water, plain mineral water or soda water instead — try these flavoured with slices of fresh lemon or lime for something different. A small amount of tea and coffee can be included during pregnancy — ask your dietitian for more information.

The Australian guidelines recommend that women who are planning a pregnancy or pregnant should not drink alcohol. For women who are breastfeeding, not drinking alcohol is the safest option.



Pregnancy weight gain

Healthy weight gain during pregnancy is important for you and your baby. It is usual to gain some weight during your pregnancy, as your baby grows. How much weight you should gain depends on your weight before you were pregnant.

Not gaining enough weight can mean that your baby may not be getting all of the nutrition they need.

Gaining too much weight during pregnancy can increase the risk of health problems such as high blood pressure, having a large baby and increased risk of birth complications, and a caesarean section. Extra weight gain can also make it more difficult to manage blood glucose levels and harder to return to a healthy weight after your baby is born.

The table below shows the recommended weight gain targets for pregnancy depending on your pre-pregnancy weight range (calculated using body mass index, or BMI, by dividing your weight in kilograms by your height in metres squared).

BMI = pre-pregnancy weight (kg) ÷ (height (m) x height (m))

Pre-pregnancy BMI	Weight range	Pregnancy weight gain (kg)	Monthly weight gain 2 nd and 3 rd trimester (kg)
< 18.5	Underweight	12.5–18	2
18.5–24.9	Healthy weight	11.5–16	1.5
25–29.9	Overweight	7–11.5	1
> 30	Obese	5–9	1

These weight gain targets do not apply to women having a multiple pregnancy, and recommendations may vary for women from different cultural backgrounds.

Have your weight checked regularly throughout your pregnancy. Discuss your individual pregnancy weight gain targets with your health professional and talk to them if you feel you are gaining too much or not enough weight.

Physical activity

When you have gestational diabetes, it is recommended that you try and be active every day.

Physical activity can help you manage your blood glucose levels and pregnancy weight gain, as well as keep you fit to prepare for the birth of your baby. It also has other benefits, such as managing pregnancy symptoms like heartburn, constipation and lower back pain.

Many types of physical activity are suitable during pregnancy, but it is important to talk to your doctor before starting or continuing any form of physical activity while you are pregnant.

For women with gestational diabetes without any other medical or pregnancy complications, aim for 30 minutes of moderate physical activity on most days of the week. This can also be broken up into shorter periods of at least 10 minutes, three times a day.

‘Moderate physical activity’ means that while being active you will have a slight but noticeable increase in breathing and heart rate (but you should still be able to hold a conversation).

Moderate activities could include:

- » swimming
- » brisk walking
- » aqua fitness classes
- » stationary cycling
- » prenatal exercise classes
- » light to moderate resistance exercise.

Pelvic floor exercises during pregnancy can also be helpful for after the baby is born. As your pregnancy progresses, you may find that some activities are more suitable than others.





You can also increase your day-to-day activity by walking to the shops, playing with your children at the park or being active around the house.

Monitoring your daily activity by keeping an exercise diary or using a device such as an activity tracker can also encourage you to be active.

To exercise safely remember to:

- » include a 5-10 minute warm up and cool down
- » drink plenty of water during and after physical activity
- » wear loose, light clothing to avoid overheating
- » avoid exercise when you are hungry, unwell or have a high temperature
- » STOP exercising and seek medical advice if you experience chest pain, dizziness, back or pelvic pain, calf pain or sudden swelling of ankles, hands or face, contractions or vaginal bleeding or a decrease in fetal movements.

During pregnancy, avoid activities that involve lying flat or increase the risk of falling, as well as contact or extreme sports.


Managing blood glucose levels

When you are active, your muscles use glucose for energy. This can help lower blood glucose levels and manage your gestational diabetes.

On days when you are less active, you may notice that your blood glucose levels will be higher, so a daily routine of physical activity can be helpful. Being active after meals (e.g. a short walk or housework) for at least 10–15 minutes can also assist with managing blood glucose levels.

Talk to your doctor or diabetes educator about the effects of exercise on your blood glucose levels, especially if you are taking insulin.

For women with gestational diabetes, moderate-intensity physical activity can help to manage blood glucose levels.



Monitoring your blood glucose levels

Blood glucose monitoring is an essential part of managing gestational diabetes.

Monitoring your own blood glucose levels will help you to:

- » better understand the effect of food and physical activity on your blood glucose levels
- » know when to seek advice from your health professionals
- » develop confidence in managing your gestational diabetes.

A diabetes educator can show you how to check your blood glucose levels using a blood glucose meter and advise you on target levels for pregnancy.

The most common times to check blood glucose levels are when you wake up in the morning (fasting) and one or two hours after each main meal. You may also be advised to check your blood glucose levels at other times.

Blood glucose targets

The following blood glucose targets are often recommended in Australia:

- » 5.0mmol/L or less before breakfast
- » 7.4mmol/L or less if you are testing one hour after the start of your meal OR
- » 6.7mmol/L or less if you are testing 2 hours after the start of your meal.

These are a general guide only and your doctor or diabetes educator will advise you on individual blood glucose targets.

Write your individual blood glucose targets in your record book/sheet.

You will also be asked to keep a record of your blood glucose readings so that your diabetes health professionals can help you look for any patterns in your blood glucose levels. They can also advise you on what to do if your blood glucose levels are outside the target range.

Your diabetes educator can help you choose a blood glucose meter. They can also help with further information on blood glucose monitoring technique, access to supplies and disposal of sharps (e.g. lancets, insulin pen needles).

Continuous glucose monitoring (CGM) is another way of checking glucose levels. However, CGM devices are not subsidised through the NDSS for gestational diabetes. This means they are more expensive than using a blood glucose meter. CGM may be useful but does not replace checking your glucose level with your blood glucose meter. Talk to your diabetes health professionals if you would like more information.

**Your doctor or
diabetes educator
will advise you on
individual blood
glucose targets.**



Medication

Some women may need medication, along with healthy eating and physical activity, to manage gestational diabetes.

Insulin

Insulin treatment may be needed to bring blood glucose levels into the target range.

Insulin is given by injection using an insulin pen device. This device can deliver the insulin at a push of a button. If you require insulin, your diabetes educator or doctor will teach you how to use the insulin device and where to inject the insulin. The injected insulin will help to lower your blood glucose level to within a range that is best for you, as well as your baby's growth and development. The insulin does not cross the placenta or affect your baby.

Your diabetes health professional will advise you of the appropriate starting dose of insulin. It is common for the insulin dose to be increased regularly throughout the second half of pregnancy (due to placental hormones causing insulin resistance). This continues until close to your delivery date.

Your diabetes team will regularly review your blood glucose levels and discuss with you the correct insulin doses to take.

Many women have worries or feel anxious about starting insulin therapy. It can take time to get used to a new treatment but your health professional is there to help you. Some women worry about giving themselves injections, but keep in mind that these days, insulin-injecting devices are relatively easy to use and have extremely fine needles. The injection of insulin will not harm your baby.

Metformin

Most diabetes tablets are not suitable for use during pregnancy, but a medication called metformin is sometimes used. Your doctor or diabetes educator will discuss whether this medication is suitable for you. If your doctor recommends metformin they will advise you of a starting dose and when to take this medication. You will need to continue to monitor your blood glucose levels and your diabetes health professionals will continue to review your doses.

Common feelings about starting medication

Some women worry that starting medication means they haven't looked after their gestational diabetes well enough or that the diabetes is getting worse. Needing medication doesn't mean that you have failed in any way, it just means that your body needs some extra help to keep blood glucose levels in the target range. Remember too, that every woman's experience with gestational diabetes and how it is managed will be different.

Every woman's experience with gestational diabetes and how it is managed will be different.



Diabetes supplies

The National Diabetes Services Scheme (NDSS) gives you access to subsidised products to help you manage gestational diabetes. These products include:

- » subsidised blood glucose testing strips
- » free insulin syringes and pen needles (if you need insulin to manage gestational diabetes)

You must be registered with the NDSS to buy subsidised products through the NDSS. Once you have your NDSS registration number or card, you can get all your NDSS products from an NDSS Access Point. Most community pharmacies are NDSS Access Points. Your diabetes health professionals can help with information about your closest Access Point or you can call the NDSS Helpline on **1800 637 700**. Medications (including insulin) are not available through the NDSS and require a prescription.

You can get a further discount on some NDSS products if you hold one of the following concession cards:

- » Health Care card
- » Pensioner concession card
- » Safety Net card
- » Department of Veterans' Affairs card



Hypoglycaemia (hypos)

If you are using insulin to manage your gestational diabetes, it is possible for your blood glucose levels to drop too low (although this is not common). A low blood glucose level is called hypoglycaemia or a 'hypo'.

This occurs when your blood glucose levels fall below 4mmol/L.

A hypo can be caused by:

- » delaying or missing a meal
- » not eating enough carbohydrate
- » being more active than usual
- » too much insulin.

Symptoms of a hypo may include nausea, weakness, trembling or shaking, sweating and dizziness. If you are taking insulin to manage gestational diabetes and feel any of these symptoms, check your blood glucose level. If it is less than 4mmol/L, treat the hypo as described below. If you can't check your blood glucose level, treat these symptoms as if you are having a hypo.

Treat hypos quickly to stop your blood glucose level from falling even lower. To treat a hypo, have some easily absorbed carbohydrate, for example:

- » Glucose gel or glucose tablets equal to 15 grams of carbohydrate OR
- » 6–7 regular size jellybeans OR
- » 3 teaspoons of sugar or honey OR
- » ½ a glass (125ml) of fruit juice OR
- » ½ a can (150ml) of regular soft drink (not 'diet').

After treating a hypo, wait 10–15 minutes then recheck your blood glucose levels to make sure they are above 4mmol/L. If symptoms continue, or if your blood glucose level is still below 4mmol/L, repeat the treatment. If your next meal is more than 20 minutes away, you will need to have some extra carbohydrate food such as a piece of fruit, glass of milk or tub of yoghurt.


Discuss managing hypos with your diabetes health professionals.

Hypos impact your ability to drive safely. If you are using insulin, discuss driving with your diabetes health professionals. They can advise you on blood glucose levels that are safe for driving and how to reduce your risk of hypos.

Always check your blood glucose before driving. Have your blood glucose monitor, hypo treatment and extra carbohydrate foods or drinks with you in the car in case of a hypo.

Do not drive a vehicle if you are having a hypo, even if you do not have symptoms. If you think you may be having a hypo while driving, pull over immediately and treat it. It is not safe to drive if you are having a hypo.

**Hypos have not been shown to cause harm to the baby.
Hypos can, however, be a risk to the safety of the mother,
so treat hypos without delay.**



The birth

Your diabetes and pregnancy health professionals will continue to monitor you and your baby throughout your pregnancy.

This will include regular checks of your blood pressure and ultrasounds to check your baby's growth and well-being. Extra blood tests will also be organised as needed. Most women with gestational diabetes will be able to deliver close to their due date and most are able to have a vaginal delivery.

If the baby grows too large or there are any other concerns about the pregnancy, the health professionals looking after your pregnancy may suggest 'inducing' the birth one or two weeks early. If an earlier birth is required, labour is usually induced after using a medication that prepares the cervix for delivery.



Birth plan

You can discuss your birth plan with your doctor or midwife. This is a good way to let them know what kind of care you would like in labour, birth and afterwards (if possible). It is a good idea to discuss your birth plan from about 36 weeks onwards, remembering that this may need to be adjusted depending on how your pregnancy is progressing.

Caesarean section

As with all pregnant women, there is a possibility that you may need a caesarean birth. Sometimes a caesarean may be required if the baby is too large or if there are other obstetric concerns such as low placenta, breech presentation or previous caesarean delivery. It is a good idea to discuss caesarean births with your health professionals so that if the need arises, you are well-prepared.

Insulin infusion (drip)

To manage blood glucose levels during labour or caesarean delivery you may need insulin by injection or by infusion (drip). This is more likely in women who have needed treatment with high doses of insulin during the pregnancy. Some women may also need a glucose drip during labour, although this is uncommon.



After the birth

After the birth, a paediatrician (a doctor who looks after babies and children), your obstetrician or a midwife will examine your baby. Your baby will be monitored carefully for the first 24–48 hours (heart rate, colour, breathing, blood glucose levels).

The midwives will perform blood glucose tests (using heel pricks) on your baby to make sure their blood glucose levels are not too low (that is, less than 2.6mmol/L). If your baby's blood glucose level is low, your baby may need to have supplementary feeds or some glucose. Talk to your midwife about using your breast milk for supplementary feeding.



Benefits of breastfeeding

Breastfeeding has many benefits, both for you and your baby. These include benefits for your baby's immune system, growth and development, and it can help with bonding between you and your baby. Breastfeeding has also been shown to have long-term health benefits for mother and baby, including reducing the risk of future type 2 diabetes.

You will be encouraged to have skin-to-skin contact with your baby and breastfeed as soon as possible after delivery. Breastfeeding at least every three hours during the first few days will help maintain your baby's blood glucose levels. Your midwife or lactation consultant can support you to establish breastfeeding and help with strategies for successful breastfeeding.

Breastfeeding information and support

For breastfeeding information and support, contact the National Breastfeeding Helpline on 1800 686 268. Support is available 24 hours a day, 7 days a week.

Local breastfeeding support networks are also available in hospitals and in the local community. Ask your midwife, lactation consultant or child and family health nurse for more information.

The practice of expressing and storing breast milk during late pregnancy is becoming more common among women with gestational diabetes, with the hope of having some colostrum (the first milk) to offer if baby has low blood glucose levels after the birth. It is very useful to learn the skill of hand expressing, so that you are prepared for when baby arrives. However, the practice of antenatal expressing may not be suitable for all women, so it is recommended that you ask your health care professionals for advice if you are considering expressing breastmilk during late pregnancy.

Medication after the birth

Medications used to treat gestational diabetes (insulin or metformin) will usually be stopped after your baby is born. Your health care team will advise you how often to monitor your blood glucose to see whether the levels have returned to normal (generally 4 to 8 mmol/L).

6–12 weeks after the birth

Most women will no longer have diabetes after the baby is born. However, some women will continue to have high blood glucose levels after delivery. An oral glucose tolerance test (OGTT) is very important to check that blood glucose levels have returned to normal. You will be advised to have this test 6–12 weeks after your baby is born (or as soon as possible after this time). An OGTT is the best way to check that your blood glucose levels have returned to the healthy range. Remember to remind your general practitioner (GP) that you have had gestational diabetes.

It is recommended that you have an oral glucose tolerance test (OGTT) 6-12 weeks after your baby is born.



Looking after your future health

Once you have had gestational diabetes, you are at a higher risk of developing type 2 diabetes in the future. If you have another pregnancy, there is also an increased risk of developing gestational diabetes again.

There are many positive steps you can take after pregnancy to reduce your risk of future diabetes. These include:

- » aiming to be in the healthy weight range
- » making healthy food choices
- » being physically active every day.

Regular checks for diabetes will be recommended every 1-3 years. These are usually a simple blood test.

Babies born to women who have had gestational diabetes also have an increased risk of childhood obesity and type 2 diabetes later in life. It is recommended that the whole family eat well and stay active to reduce this risk.

After the birth of your baby, you will be sent another booklet, *Life after gestational diabetes*, as well as regular reminders for follow-up type 2 diabetes checks and healthy lifestyle information.

About half of all women who have had gestational diabetes will develop pre-diabetes or type 2 diabetes within 10-20 years.



National Diabetes Services Scheme

What is the NDSS?

The NDSS supports people with diabetes by giving them access to reliable and affordable services and products.

The NDSS is an initiative of the Australian Government. Diabetes Australia has administered the NDSS on behalf of the Australian Government since its inception in 1987. Registration with the NDSS is free and open to all Australians who have been diagnosed with diabetes and have a valid Medicare card.

The NDSS provides a range of services and support to help you manage your diabetes. These include the NDSS Helpline on **1800 637 700** for advice on diabetes management, subsidised NDSS products, and information, support and education to help you learn more about managing your diabetes.

Where can I access NDSS services and products?

You can access NDSS services and support by calling the NDSS Helpline on **1800 637 700**. The NDSS Helpline can also put you in contact with your NDSS Agent. NDSS Agents are the diabetes organisations in each state and territory, and the names and contact details of all Agents are provided at the end of this book.

You can access subsidised NDSS products in all states and territories through NDSS Access Points, which are usually community pharmacies. You can find your closest Access Point by calling the NDSS Helpline on **1800 637 700**.

Services and support

Through the NDSS people with gestational diabetes can access:

- » Support services such as online self-management education programs for practical help and guidance
- » Digital diabetes information and resources
- » Lower cost diabetes products.

National Gestational Diabetes Register

The National Gestational Diabetes Register was established within the NDSS to help women who have had gestational diabetes manage their health during pregnancy and into the future. When you first register with the NDSS with gestational diabetes you are automatically included on the Register. As part of the Register, you will be sent reminders about having regular type 2 diabetes checks after your baby is born.

If you would like more information about the National Gestational Diabetes Register, go to ndss.com.au or call the NDSS Helpline on **1800 637 700**.

The logo for the National Diabetes Services Scheme (NDSS) features the letters 'nDSS' in a bold, blue, sans-serif font. The 'n' is lowercase, while 'DSS' is uppercase.

National Diabetes Services Scheme

An Australian Government Initiative

NDSS agent contact details

New South Wales and the Australian Capital Territory

Diabetes Australia

diabetesaustralia.com.au

Northern Territory

Healthy Living NT

healthylivingnt.org.au

Queensland

Diabetes Australia

diabetesaustralia.com.au

South Australia

Diabetes SA

diabetessa.com.au

Tasmania

Diabetes Australia

diabetesaustralia.com.au

Victoria

Diabetes Victoria

diabetesvic.org.au

Western Australia

Diabetes WA

diabeteswa.com.au

Acknowledgments:

The Australian Government and Diabetes Australia wish to acknowledge the valuable contribution and support of the National Diabetes Services Scheme (NDSS) Diabetes in Pregnancy Expert Reference Group (ERG) who provided content expertise in the update of this publication. 2019 ERG members include – Associate Professor Glynis Ross (chair), Associate Professor Alison Nankervis, Associate Professor Ralph Audehm, Dr Christel Hendrieckx, Alison Barry, Dr Cindy Porter, Dr Melinda Morrison and Renza Scibilia.

The input of the dietitians who provided expertise in the update of the nutrition content of this booklet is acknowledged and appreciated. Thank you to Julia Zinga, Robyn Barnes, Dr Susan de Jersey, Dr Shelley Wilkinson, Rachel Hayes, Judith Ingle, Anita Marshall, Elin Donaldson, Laura Barsha, Effie Houvardas and members of the National Diabetes Australia Dietitians group.

Diabetes Australia is grateful for the assistance of consumers and health professionals who provided feedback and suggestions during the review and update of this booklet. This booklet was modified in collaboration with The Australian Centre for Behavioural Research in Diabetes, a partnership for better health between Diabetes Victoria and Deakin University.

For further information regarding this publication, its development or availability call the NDSS Helpline on **1800 637 700** or email **info@ndss.com.au**

